

COSMIC ACCELERATION AND M-THEORY¹

Paul K. Townsend

Department of Applied Mathematics and Theoretical Physics,
Centre for Mathematical Sciences, University of Cambridge,
Wilberforce Road, Cambridge CB3 0WA, UK

ABSTRACT

The status of accelerating four-dimensional universes obtained by time-dependent compactifications of 10 or 11 dimensional supergravity is reviewed, as is the ‘no-go’ theorem that they evade. All flat cosmologies for a simple exponential potential are found explicitly. It is noted that transient acceleration is generic, and unavoidable for ‘flux’ compactifications. Included is an eternally accelerating flat cosmology without a future event horizon.

¹To appear in the proceedings of ICMP2003, Lisbon, Portugal, July 2003

The current consensus that the Universe is undergoing accelerated expansion presents a challenge to the other current consensus that cosmology should be derivable from String/M-theory, which has 10 or 11-dimensional supergravity as its low-energy effective field theory. There are two aspects to this challenge. One arises from the fact that there is no known formulation of String/M-theory in a spacetime with a future cosmological event horizon [1, 2], whereas this is a typical feature of universes undergoing late time eternal acceleration. Of course, we don't know whether the accelerated expansion of the Universe will continue forever, so there is no real conflict with observations here. Still, it remains to find compactifications of String/M-theory for which the effective 4-dimensional theory admits a homogeneous and isotropic (FLRW) spacetime undergoing accelerated expansion, and the difficulty of finding such compactifications is the other aspect of the challenge posed by cosmic acceleration. Note that *two* periods of acceleration are required to explain both inflation in the early universe and acceleration in the current cosmological epoch.

To see what cosmic acceleration entails, consider, a 4-dimensional FLRW spacetime in standard coordinates. The metric is

$$ds^2 = -dt^2 + S^2(t) \left[(1 - kr^2)^{-1} dr^2 + r^2 d\Omega_2^2 \right], \quad (1)$$

where $S(t)$ is the scale factor and $k = -1, 0, 1$ depending on whether the universe is open, flat or closed, and $d\Omega_2^2$ is the $SO(3)$ invariant metric on the unit 2-sphere. A computation of the Ricci tensor shows that

$$R_{00} = -\partial_t^2 S. \quad (2)$$

It follows that an accelerating universe requires $R_{00} < 0$. However, the Einstein field equations imply that

$$R_{00} = (T_{00} + g^{ij}T_{ij}) \quad (3)$$

and the Strong Energy Condition (SEC) on the matter stress tensor requires the right hand side to be non-negative. Thus, accelerated expansion is possible in a universe governed by Einstein's equations only if the matter in it violates the SEC.

From a purely 4-dimensional perspective the fundamental condition on the matter stress tensor is the Dominant Energy Condition (DEC), which requires $p \geq -\rho$ for a perfect fluid of pressure p and energy density ρ . In contrast, the SEC requires only that $p \geq -\frac{1}{3}\rho$, and is typically violated in theories with a positive scalar potential V . For example, given a single scalar field ϕ and the Lagrangian density

$$\mathcal{L} = \sqrt{-\det g} \left[\frac{1}{4}R - \frac{1}{2}(\partial\phi)^2 - V \right], \quad (4)$$

the Einstein equations imply that

$$R_{00} = 2[(\partial_t\phi)^2 - V]. \quad (5)$$

If $V > 0$ then R_{00} will be negative whenever $\partial_t\phi = 0$, implying an accelerating universe. There will always be *some* cosmological solutions for which $\partial_t\phi$ passes

through zero. In fact, as will become clear in due course, such solutions are the rule rather than the exception! Thus, all one needs to get an accelerating universe from String/M-theory is a compactification for which the effective 4-dimensional theory has a positive scalar potential V , or at least one that is positive in some region of the space of scalar fields.

Although the SEC is in no way fundamental, in the sense that its violation would not imply a violation of fundamental physical principles, it is satisfied by the stress tensor of the D=10 and D=11 supergravities that serve as the low energy effective theories of String/M-theory². This fact has consequences for the potential V in the effective 4-dimensional theories that result from String/M-theory compactifications, as first pointed out in a 1985 article of Gibbons [3], and more recently by Maldacena and Nuñez [4]. I will now summarize this ‘no-go theorem’. Consider a D-dimensional spacetime with metric

$$ds_D^2 = \Omega^2(y) ds_4^2(x) + ds_n^2(y) \quad (n = D - 4) \quad (6)$$

where ds_4^2 is the metric of some 4-dimensional spacetime (with coordinates x), and ds_n^2 is the metric of some compact non-singular n -manifold $\mathcal{M}$ (with coordinates y). The non-vanishing function $\Omega(y)$ is a ‘warp factor’. A calculation shows that

$$R_{00}^{(D)}(x, y) = R_{00}(x) - \frac{1}{4} \Omega^{-2}(y) \nabla_y^2 \Omega^4(y). \quad (7)$$

Multiplying by Ω^2 and integrating over $\mathcal{M}$ we deduce that

$$\left[\int_{\mathcal{M}} \Omega^2 \right] R_{00} = \int_{\mathcal{M}} \Omega^2 R_{00}^{(D)}, \quad (8)$$

and hence that $R_{00} \geq 0$ if $R_{00}^{(D)} \geq 0$.

This result might appear to rule out the possibility of an accelerating universe arising from compactification of 10 or 11-dimensional supergravity. In view of our earlier remark that acceleration is always possible when the 4-dimensional scalar potential V is positive, this would be equivalent to the statement that the potential arising from such compactifications is never positive, a statement that is false, and was known to be false well before the no-go theorem was formulated. In fact³, all that can be inferred about the potential V is that *it has no stationary points with $V > 0$* . To see that a positive potential without a stationary point is not excluded, it suffices to note that the field equations imply, under the given circumstances, that at least one scalar field is time-dependent. This scalar field could be one that arises from the mode expansion on the compact manifold $\mathcal{M}$, in which case the metric on $\mathcal{M}$ will be time-dependent. But the theorem assumed time-independent $\mathcal{M}$, and is therefore not applicable. One can see from this that the no-go theorem

²The ‘massive’ IIA supergravity is an exception but, for a reason to be explained later, this exception makes no difference.

³This point was made in [5], in a note added to the published version; it has also been made, independently, in [6].

is actually a very *weak* constraint on the positive potentials that might, in principle, arise from compactification of String/M-theory! There are many potentials that it would allow but which, nevertheless, do not seem to be obtainable. For example, for many compactifications there is a consistent truncation to a single scalar field ϕ with a potential of the form

$$V = \Lambda e^{-2a\phi} \quad (\Lambda > 0) \quad (9)$$

for (dilaton coupling) constant a (which we may assume to be positive). Any value of a would be permitted by the no-go theorem but only $a > 1$ arises in practice. This might be expected on the grounds that $a < 1$ allows an eternally accelerating cosmology with a future event horizon. This suggests the conjecture that such (Einstein conformal frame) spacetimes cannot arise from classical compactification of String/M-theory; if true (there is no known counterexample) this would impose much stronger constraints on the potential V than the no-go theorem. In particular, it would exclude $a < 1$ (but not $a = 1$, as will be shown later).

Let us now turn to the question of how one gets positive potentials from (classical) compactification of higher-dimensional theories satisfying the SEC. These arise in one of two ways:

- **Flux compactifications:** in this case a positive potential is generated by non-zero flux of antisymmetric tensor fields. The prototype is the T^7 compactification of 11-dimensional supergravity with non-vanishing 4-form field strength [7], which yields an exponential potential of the form (9) with⁴ $a = \sqrt{7}$, the scalar field arising from the ‘breathing mode’ of T^7 . The 4-form is dual to a 7-form proportional to the volume form of T^7 ; more generally, some k -form field strength will be set equal to a closed but not exact k -form on the compact space $\mathcal{M}$. Flux compactifications seem only to yield ‘steep’ exponential potentials with $a \geq \sqrt{3}$.
- **Hyperbolic compactifications:** in this case the compact space is a space of constant negative curvature. The fact that hyperbolic compactifications produce a positive potential was observed by Bremer et al. [9], and they were investigated by Kehagias and Russo [10] in the context of String/M-theory. Several attractive features (for example, the absence of moduli other than the volume) were noted and exploited in a cosmological context by Kaloper et al. [11], and their possible relevance to cosmic acceleration was noted by Wohlfarth and the author [5]. One could consider the prototype to be the compactification of 11-dimensional supergravity on a 7-dimensional compact hyperbolic space. In this case one finds a potential for the breathing-mode scalar ϕ of the form (9) with $a = 3/\sqrt{7}$. In general, hyperbolic compactifications seem always lead to ‘gentle’ exponential potentials with $1 < a < \sqrt{3}$.

⁴This value, which corresponds to $\Delta = 4$ in the notation of [8] was given incorrectly in that paper. The correct value is the same as the one found from T^6 compactification of the massive IIA supergravity, which is why the existence of this SEC violating theory does not alter our conclusions.

In general, a positive multi-scalar potential can be generated by a combination of both mechanisms, in which case it takes the form of a sum of products of exponentials of canonically normalized scalar fields. However, the simple case of a single exponential for a single scalar field is sufficient for an understanding of the physics and here we shall consider only this case.

The qualitative features of cosmologies derived from (4) with a potential of the form (9) were analysed in a 1987 paper of Halliwell [12], although the fact that there is typically a period of transient acceleration in the $a > 1$ cases was not noticed there. In 2002 Cornalba and Costa [13] noted the existence of a period of acceleration in a $k = -1$ cosmology arising from a flux compactification⁵. More recently, an explicit time-dependent hyperbolic compactification of the vacuum Einstein equations was shown to yield an Einstein-frame $k = 0$ universe in which a decelerating epoch with

$$S \sim t^{\frac{1}{3}}, \quad e^{\phi} \sim t^{-\frac{1}{\sqrt{3}}}, \quad (10)$$

is followed by a period of transient acceleration [5]. This solution was subsequently shown to be the vanishing flux limit of a rather general class of solutions of Einstein's equations known as S-branes, and the phenomenon of transient acceleration was found to be a generic feature of these solutions [14]. As observed by Emparan and Garriga [15], this is an immediate corollary of the positive potentials generated by flux and hyperbolic compactifications. Consider the simple case of an exponential potential of the form (9) with $a > 1$. The initial conditions implied by (10) are $\phi \gg 1$ with $\dot{\phi} < 0$. Any such cosmological solution can be viewed as a ball rolling, with friction, up the potential. Clearly, it must reach a maximum at which $\dot{\phi} = 0$ and at this point the expansion of the universe is accelerating, for the reason explained previously. Subsequently, the ball starts to roll back down the hill; the late-time behaviour will depend on the value of a and also on k , but in all cases the universe will be decelerating. For example, for $a^2 < 3$ and $k = 0$ the late-time behaviour will be given by the power-law $k=0$ attractor solution

$$S \sim t^{1/a^2}, \quad e^{\phi} \sim t^{1/a}. \quad (11)$$

Nearby trajectories with $k \neq 0$ will eventually approach a Milne universe attractor or collapse to a big crunch singularity. Note that the compact Kaluza-Klein space $\mathcal{M}$ starts at infinite volume and ends at infinite volume; the acceleration of the 4-dimensional cosmology is associated to a ‘bounce’ of the compact space off its minimal volume.

The above explanation of the period of transient acceleration relies only on the positivity of the potential V and makes no distinction between flux compactifications and hyperbolic compactifications. However, Halliwell's analysis [12] shows that just as the ‘critical’ value $a = 1$ separates qualitatively different behaviours of cosmological

⁵The acceleration was claimed to occur in the neighbourhood of a resolved cosmological singularity. However, for reasons that will hopefully become clear below, the accelerating epoch is necessarily far from the cosmological singularity.

trajectories in the class of models under discussion, so does the ‘hyper-critical’ value $a = \sqrt{3}$, which also separates hyperbolic from flux compactifications. To see this, we introduce a new time parameter τ such that

$$d\tau = e^{-a\phi} dt, \quad (12)$$

and set $S = e^{\alpha(\tau)}$. Letting an overdot indicate differentiation with respect to τ , we find that the ϕ equation of motion is

$$\ddot{\phi} - a\dot{\phi}^2 + 3\dot{\alpha}\dot{\phi} = 2a, \quad (13)$$

while the Friedmann equation is

$$\dot{\alpha}^2 - \frac{1}{3}\dot{\phi}^2 = \frac{2}{3} - \frac{ke^{2a\phi}}{S^2}. \quad (14)$$

For $k = 0$, the above two equations are equivalent to

$$\ddot{\phi} = 3\dot{\alpha} (a\dot{\alpha} - \dot{\phi}), \quad (15)$$

and

$$3\dot{\alpha}^2 - \dot{\phi}^2 = 2, \quad (16)$$

which is a hyperbola separating the $k = -1$ and $k = +1$ trajectories. The $\dot{\alpha} > 0$ branch of the hyperbola corresponds to an expanding universe. We can parametrize this branch by writing

$$\dot{\alpha} = \frac{1}{\sqrt{6}} (\xi + \xi^{-1}), \quad \dot{\phi} = \frac{1}{\sqrt{2}} (\xi - \xi^{-1}), \quad (\xi > 0). \quad (17)$$

Equation (15) then becomes

$$\dot{\xi} = \frac{1}{\sqrt{2}} \left[(a + \sqrt{3}) + (a - \sqrt{3}) \xi^2 \right]. \quad (18)$$

We now see why $a = \sqrt{3}$ is special. For $a < \sqrt{3}$ there is a fixed point solution

$$\xi = \xi_0 \equiv \left(\frac{\sqrt{3} + a}{\sqrt{3} - a} \right)^{\frac{1}{2}}, \quad (19)$$

which is just the power-law attractor solution (11). The fixed point separates two other $k = 0$ trajectories:

$$(i) \ \xi = \xi_0 \coth \gamma\tau \quad (ii) \ \xi = \xi_0 \tanh \gamma\tau, \quad \gamma = \left(\frac{3 - a^2}{2} \right)^{\frac{1}{2}} \quad (20)$$

where $\tau > 0$. Only case (ii) includes $\xi = 1$, and hence $\dot{\phi} = 0$; this solution undergoes a period of acceleration whereas the other does not. In either case one can integrate (17). In case (i) one has

$$S^{\sqrt{3}} \propto (\cosh \gamma\tau)^{\lambda_+} (\sinh \gamma\tau)^{\lambda_-}, \quad e^{\phi} \propto (\cosh \gamma\tau)^{-\lambda_+} (\sinh \gamma\tau)^{\lambda_-}, \quad (21)$$

where

$$\lambda_{\pm} = \frac{1}{\sqrt{3} \pm a}. \quad (22)$$

There is a big-bang singularity at $\tau = 0$, near which

$$S \sim t^{\frac{1}{3}}, \quad e^{\phi} \sim t^{\frac{1}{\sqrt{3}}}, \quad (23)$$

so the volume of the compact Kaluza-Klein space $\mathcal{M}$ is initially zero. Subsequently the solution approaches the attractor (11). In case (ii) we have

$$S^{\sqrt{3}} \propto (\cosh \gamma\tau)^{\lambda_-} (\sinh \gamma\tau)^{\lambda_+}, \quad e^{\phi} \propto (\cosh \gamma\tau)^{\lambda_-} (\sinh \gamma\tau)^{-\lambda_+}. \quad (24)$$

This behaves initially as in (10) but then passes through a period of acceleration before approaching the attractor (11).

Both the above solutions were found in [5], for particular values of a in the range $1 < a < \sqrt{3}$, as solutions of the $D \geq 6$ vacuum Einstein equations with a compact hyperbolic $D-4$ dimensional manifold of time-dependent volume. As solutions of the 4-dimensional effective theory with Lagrangian density (4), they are actually valid for $0 \leq a < \sqrt{3}$, in particular for $a = 1$. The power-law attractor solution in this case has $S \sim t$ and hence zero acceleration, so the case (ii) solution that approaches it asymptotically must be *eternally* accelerating⁶. In fact, the late time behaviour is

$$e^{\phi} \sim t + \frac{2\sqrt{3}}{t} + \mathcal{O}(t^{-2}), \quad S \sim t + \frac{4}{\sqrt{3}t} + \mathcal{O}(t^{-2}). \quad (25)$$

from which one sees that $\partial_t^2 S > 0$. One might suspect from this fact that there would be a future cosmological event horizon, in which case $a = 1$ would be excluded by the (admittedly conjectural) stronger form of the no-go theorem proposed earlier in this article. However, it has been shown by Boya et al. [17] that if the acceleration tends to zero asymptotically, as it does in this case, then there is no cosmological event horizon.

Halliwel's qualitative analysis of all cosmological trajectories can similarly be made quantitative for $k = 0$ when $a \geq 3$. Consider first the $a > \sqrt{3}$ case. The solution of (18) is

$$\xi = \left(\frac{a + \sqrt{3}}{a - \sqrt{3}} \right)^{\frac{1}{2}} \tan \omega\tau, \quad \omega = \left(\frac{a^2 - 3}{2} \right)^{\frac{1}{2}}, \quad (26)$$

where $0 < \tau < \pi/2$. The equations (17) can now be integrated to yield

$$S^{\sqrt{3}} \propto (\cos \omega\tau)^{\lambda_-} (\sin \omega\tau)^{\lambda_+}, \quad e^{\phi} \propto (\cos \omega\tau)^{\lambda_-} (\sin \omega\tau)^{-\lambda_+}. \quad (27)$$

The asymptotic behaviour as $\log t \rightarrow \pm\infty$ is

$$S \sim t^{\frac{1}{3}}, \quad e^{\phi} \sim t^{\pm 1/\sqrt{3}}. \quad (28)$$

⁶A similar observation was made in [16] in the $a > 1$ case for trajectories that approach the $k = -1$ Milne attractor.

In between there is a period of acceleration. For $a = \sqrt{3}$ we have simply $\xi = \sqrt{6}\tau$ and hence

$$S^3 \sim \tau^{\frac{1}{2}} e^{\frac{3}{2}\tau^2}, \quad e^{\sqrt{3}\phi} \propto \tau^{-\frac{1}{2}} e^{\frac{3}{2}\tau^2}. \quad (29)$$

The late time behaviour now involves logarithmic corrections to the power law behaviour of the $a > \sqrt{3}$ case.

Despite the differences between flux compactifications and hyperbolic compactifications, all the above cases of accelerating $k = 0$ universes are qualitatively similar. There is always a big-bang singularity near which the scale factor behaves as in (10). Inspection of the phase portraits in [12] shows that this is also true for the $k \neq 0$ cosmologies. This is not surprising because the singularity theorems that guarantee a cosmological singularity rely on the SEC which is violated by the 4-dimensional effective theory only in a *later* epoch. In addition, the acceleration leads to only a few e-foldings, insufficient for any application to inflation in the early universe; this disappointing conclusion is confirmed by systematic analyses of the possibilities of hyperbolic-flux compactifications involving many scalar fields [16, 18]. However, the mechanism may be relevant to acceleration in the current cosmological epoch [19]. One of its attractive features is that any time-dependence of four-dimensional ‘constants’ due to the time-dependence of the compact space is absent precisely during the period of acceleration, i.e. now!

Implicit in everything discussed so far in this article is the assumption that String/M-theory is adequately described for cosmological purposes by the classical effective 10 or 11 dimensional supergravity theories. It seemed worthwhile to fully explore the implications of this assumption, but it now also seems that it must be discarded because although it has been possible to find models with transient acceleration of possible relevance to the ‘observed’ acceleration of the current cosmological epoch, it has not been possible to find models that allow a sufficient period of early universe inflation. This is not a disaster; String/M-theory includes orientifolds that violate (at least locally) the SEC [20], and branes which can produce brane-instanton quantum corrections to the potential V . It has been shown [21] that when these effects are taken into account it is possible for the potential V to have a local minimum that could lead to inflation⁷. Nevertheless, although string and brane effects might yield potentials V that are quite different from those obtainable from the classical compactifications of effective supergravity theories considered here, some of the lessons learned from the latter case may prove valuable. If one re-interprets a in (15) and (16) to be shorthand for $V'/2V$ then these equations describe the $k = 0$ cosmologies for *any* positive potential V . As long as $a \geq \sqrt{3}$, there will be no fixed point on the $k = 0$ hyperbola and hence a single $k = 0$ trajectory, which will necessarily include a period of acceleration. Thus, for a large class of potentials, including all those that arise classically from ‘product’ flux compactifications, an accelerating epoch is not only possible, but (assuming a flat universe) unavoidable! If $a < \sqrt{3}$ at some point

⁷Such a potential would allow classical solutions with a future cosmological event horizon, but since quantum mechanics has now been invoked one can presumably invoke it again to argue that the universe will eventually tunnel out of the metastable vacuum.

then fixed points will occur, and acceleration can be avoided. But note that we are now discussing how to avoid acceleration rather than how to achieve it! When it was noted in [5] that hyperbolic compactification can yield both a flat 4-dimensional universe that undergoes a period of acceleration *and* a flat 4-dimensional universe that is always decelerating, it was the former case that appeared remarkable, but actually it is the latter case, an eternally decelerating universe, that is exceptional. Of course, the acceleration is transient, so even if it is generic one could ask why we happen to be around to observe it. But is it really any more surprising that we live at an atypical time in a typical universe than that we live at a typical time in a atypical universe?

Acknowledgements. The author thanks Michael Douglas, Gary Gibbons, Jaume Garriga, Jaume Gomis, Shamit Kachru and Mattias Wohlfarth for helpful discussions.

References

- [1] T. Banks, *Some thoughts on the quantum theory of de Sitter space*, astro-ph/0305037.
- [2] S. Hellerman, N. Kaloper and L. Susskind, *String theory and quintessence*, JHEP 0106:003,2001; W. Fischler, A. Kashani-Poor, R. McNees and S. Pabhan, *The acceleration of the universe, a challenge for string theory*, JHEP 0107:003,2001.
- [3] G.W. Gibbons, *Aspects of supergravity theories in Supergravity and related topics*, eds. F. de Aguila, J.A. de Azcaárraga and L. Ibañez, (World Scientific, 1985).
- [4] J. Maldacena and C. Nuñez, *Supergravity description of field theories on curved manifolds, and a no-go theorem*, J. Mod. Phys. **A 16** (2001) 822.
- [5] P.K. Townsend and M.N.R. Wohlfarth, *Accelerating cosmologies from compactification*, Phys. Rev. Lett. **91**, (2003) 061302.
- [6] S. de Alwis, *On potentials from fluxes*, hep-th/0307084.
- [7] A. Aurilia, H. Nicolai and P.K. Townsend, *Hidden constants: The theta parameter of QCD and the cosmological constant of N=8 supergravity*, Nucl. Phys. **B176** (1980) 509.
- [8] P.K. Townsend, *Quintessence from M-theory*, JHEP 0111:042,2001.
- [9] M.S. Bremer, M.J. Duff, H. Lü, C.N. Pope and K.S. Stelle, *Instanton cosmology and domain walls from M-theory and string theory*, Nucl. Phys. **B543** (1999) 321.

- [10] A. Kehagias and J.G. Russo, *Hyperbolic spaces in String and M Theory*, JHEP 0007:027,2000.
- [11] N. Kaloper, J. March-Russell, G.D. Starkman, M. Trodden, *Compact hyperbolic extra dimensions: Branes, Kaluza-Klein modes and cosmology*, Phys. Rev. Lett. **85** (2000) 928.
- [12] J.J. Halliwell, *Scalar fields in cosmology with an exponential potential*, Phys. Lett. **B185** (1987) 341.
- [13] L. Cornalba and M.S. Costa, *A new cosmological scenario in String Theory*, Phys. Rev. **D 66** (2002) 066001.
- [14] N. Ohta, *Accelerating cosmologies from S-branes*, Phys. Rev. Lett. **91** (2003) 061303; S. Roy, *Accelerating cosmologies from M/String-theory compactifications*, Phys. Lett. **B567** (2003) 322; M.N.R. Wohlfarth, *Accelerating cosmologies and a phase transition in M-theory*, Phys. Lett. **B563** (2003) 1; N. Ohta, *A study of accelerating cosmologies from Superstring/M-theory*, hep-th/0304172; C-M. Chen, P-M Ho, I.P. Neupane and J.E. Wang, *A note on acceleration from product space compactification*, JHEP 0307:017,2003.
- [15] R. Emparan and J. Garriga, *A note on accelerating cosmologies from compactifications and S-branes*, JHEP 0305:028,2003.
- [16] C-M Chen, P-M Ho, I.P. Neupane, N. Ohta and J.E. Wang, *Hyperbolic space cosmologies*, hep-th/0306291.
- [17] L.J. Boya, M.A. Per and A.J. Segui, *Graphical and kinematical approach to cosmological horizons*, Phys. Rev. **D 66**, 064009 (2002).
- [18] M.N.R. Wohlfarth, *Inflationary cosmologies from compactification?*, hep-th/0307179.
- [19] M. Gutperle, R. Kallosh and A. Linde, *M/String theory, S-branes and accelerating universe*, JCAP 0307:001,2003.
- [20] S.B. Giddings, S. Kachru and J. Polchinski, *Heirarchies from fluxes in string compactifications*, Phys. Rev. **D 66** (2002) 106006.
- [21] S. Kachru, R. Kallosh, A. Linde and S.P. Trivedi, *De Sitter vacua in string theory*, Phys. Rev. **D68** (2003) 046005; S. Kachru, R. Kallosh, A. Linde, J. Maldacena, L. McAllister and S.P. Trivedi, *Towards inflation in string theory*, hep-th/0308055.